

POST - OPERATIVE INSTRUCTIONS

Pain Management

- During the procedure local anaesthetic is injected into the operative site which should numb the area for 6-12 hours after your procedure.
- Upon discharge you may begin to feel some discomfort for which you may take:
 - Panadol/Panadeine/Panadeine Forte, 2 tablets every 6 hours regularly for the first 48-72 hours.
 - Anti-inflammatory e.g. Nurofen can be used in addition if required.
 - Endone may be prescribed for more extensive procedures. Take this for break through pain.
- If your pain is not controlled with the above measure contact WPRS for advice.

Wound Care

- If you have a dressing in place then please keep dry and intact, unless you are informed otherwise. Certain dressings (mefix and tegaderm) can be wet and patted dry. If your dressing falls off, do not be concerned as it will not harm the operative site. You should contact WPRS during office hours for advice regarding the dressing.
- If any bleeding occurs upon discharge, please apply pressure to the site for 10 minutes. If bleeding continues, repeat the above. If bleeding persists contact WPRS.
- Small amounts of ooze from your wound is normal. Please consult WPRS if you are concerned by the amount or type of discharge.
- Bruising and minimal swelling can be expected around the wound, but this should remain stable and improve in following days.

Activity

- Gentle home activity is recommended for the first 2-4 weeks.
- Avoid heavy lifting, strenuous exercise or activity for 6 weeks.
- 2-4 weeks following your surgery you can consider returning to driving, if you feel comfortable and safe to do so.
- Exercise your ankles, calves whilst in your recovery to minimise your risk for deep vein thrombosis.

Diet

- After most abdominal surgeries, the gut experiences some form of paralysis (laziness). This manifest in the form of bloating, nausea and/or vomiting. Pain medication can contribute to this, therefore use pain medication as prescribed with a high fibre diet.
- Gut motility/activity also differs from person to person. For this reason, we cannot give strict guidelines.
- Normally start with clear fluids after surgery or the next day. We progress to full fluid/light diet in the following day or 2.
- Day 2-3 As your gut starts to work (rumble) and you have bowel actions, you can gradually pace yourself back to your normal diet. This doesn't mean you can eat what you want when home. You will still need to be gentle on the gut and take it easy and slow with the reintroduction of oral intake.

Follow-up

- Day 7-10 post procedure you will have a post-operative appointment at WPRS.
- WPRS staff will have attached an appointment card to this handout advising you of this. If you haven't received a card or appointment, please contact the WPRS rooms on (03) 5562 5330.

If you have any concerns please contact WPRS on (03) 5562 5330.

- Within clinic hours your call will be transferred to clinical nurse and managed appropriately.
- After hours you may leave a message or your call will be diverted to the on-call Surgeon.

In the unlikely event you are unable to contact staff at WPRS please call St John of God Hospital, Warrnambool on (03) 5564 0600.